CUMULATIVE AUTHOR INDEX
Annals of Entomology
Volume 98, 2005

A

Ahmad, I., 282
Akhmetiev, M. A., 503
Alarie, Y., 150, 417
Almeida, R.P.P., 755, 775
Aluja, M., 252
Alyokhin, A., 989
Ammar, E.-D., 820
Archibald, S. B., 503
Arciniega, O., 960
Austin, J. W., 273, 395, 980

B

Backus, E. A., 787
Bagnères, A.-G., 395
Baker, M. B., 989
Bakri, A., 1
Barr, N. B., 173
Barrera, E., 48
Baselga, A., 558, 896
Bashan, M., 575
Bayer, M. A., 608
Beanland, L., 756
Benavides, P., 359
Bertone, M., 309
Bextine, B. R., 814
Billick, I., 491
Bilton, D. T., 417
Blanc, S., 763
Blua, M. J., 775
Boonchu, N., 113
Bostanian, N. J., 565
Bouchard, P., 565
Bousquets, J. L., 227
Brailovsky, H., 48
Brannan, M., 601
Briano, J. A., 85
Brower, A.V.Z., 873, 1002
Brower, L. P., 682
Brown, B. V., 55
Brown, C. G., 711
Brown, J. K., 827
Brown, R., 491
Browne, J., 951
Brust, M. L., 973
Bruton, B., 770
Bustillo, A. E., 359

C

Cabrera, N., 643
Caceres, C., 119
Caesar, R. M., 931
Cakmak, O., 575
Calcaterra, L. A., 85
Calvo, D., 191
Candan, S., 346
Carew, M. E., 587
Carpintero, D. L., 694
Carstens, B. C., 165
Chaiwong, T., 113
Chattrukul, U., 887
Chen, B.-H., 475
Chi, H., 475
Chi, Y., 470
Choate, B. A., 496
Choi, W.-Y., 636
Cisneros, R., 960
Cognato, A. I., 931
Conn, J. E., 908
Cordo, H. A., 66
Coşkun, M., 336
Cui, L., 173

D

Dabert, J., 13
D'Adamo, P., 236
Danley, P. D., 219
Dastur, S. R., 989
Davis, A. K., 682
Delprat, A., 119
DeWalt, R. E., 941
Díaz, E., 960
Diez-Rodríguez, G. I , 409
Disney, H., 113
Dorado, J., 643
Dronnet, S., 395
Duyck, P. F., 402

E

Ehrnsberger, R., 13
Ellersieck, M., 787
Emre, I., 336
Engel, M. S., 73
Evans, W. G., 738

F

Fang, J., 351
Favret, C., 941
Fereres, A., 763
Ferris, I., 1
Ferro, D. N., 989
Filippi, L., 134, 143
Flakus, A., 108
Fletcher, J., 770
Fontal-Cazalla, F., 63
Forister, M. L., 295
Foster, P. G., 918
Franz, G., 119
Fuller, S., 126
Funk, D. J., 711

G

Galun, R., 302
Galvão, C., 861
Garofalo, R. S., 329
Garrido, A., 747
Gera, A., 351
Gillette, N., 931
Godoy, C., 444
Goodpasture, C., 227
Goolsby, J. H., 887
Goughnour, R. B., 703
Goulet, H., 565
Graves, W. R., 690, 926
Green, J., 309
Griswold, M. W., 673
Gut, L. J., 580

H

Habibi, J., 787
Hanski, I., 382
Harshman, D. B., 814
Hazel, W., 996
Heather, N., 1
Henderson, G., 340, 732
Hendrichs, J., 1
Hill, J. M., 552
Hironaka, M., 134, 143
Hirose, E., 515
Hirose, Y., 291
Hoagstrom, C. W., 241
Hoback, W. W., 973
Hoffmann, A. A., 587
Hogenhout, S. A., 820
Honda, J. Y., 291
Horton, D. R., 527, 608
Hoy, C. W., 756
Hussein, S., 227

I

Idris, A. M., 827
Infante, F., 483
Israely, N., 77, 302
Iwakuma, S., 143

J

Jacas, J.-A., 747
Jackson, B. C., 814
Jameson, M. L., 658
Johnson, D. G., 165
Jones, W. A., 322
Julien, M., 643
Jurberg, J., 861

K

Kankare, M., 382
Keena, M. A., 536

CUMULATIVE AUTHOR INDEX
Annals of Entomology
Volume 98, 2005

A

Ahmad, I., 282
Akhmetiev, M. A., 503
Alarie, Y., 150, 417
Almeida, R.P.P., 755, 775
Aluja, M., 252
Alyokhin, A., 989
Ammar, E.-D., 820
Archibald, S. B., 503
Arciniega, O., 960
Austin, J. W., 273, 395, 980

B

Backus, E. A., 787
Bagnères, A.-G., 395
Baker, M. B., 989
Bakri, A., 1
Barr, N. B., 173
Barrera, E., 48
Baselga, A., 558, 896
Bashan, M., 575
Bayer, M. A., 608
Beanland, L., 756
Benavides, P., 359
Bertone, M., 309
Bextine, B. R., 814
Billick, I., 491
Bilton, D. T., 417
Blanc, S., 763
Blua, M. J., 775
Boonchu, N., 113
Bostanian, N. J., 565
Bouchard, P., 565
Bousquets, J. L., 227
Brailovsky, H., 48
Brannan, M., 601
Briano, J. A., 85
Brower, A.V.Z., 873, 1002
Brower, L. P., 682
Brown, B. V., 55
Brown, C. G., 711
Brown, J. K., 827
Brown, R., 491
Browne, J., 951
Brust, M. L., 973
Bruton, B., 770
Bustillo, A. E., 359

C

Cabrera, N., 643
Caceres, C., 119
Caesar, R. M., 931
Cakmak, O., 575
Calcaterra, L. A., 85
Calvo, D., 191
Candan, S., 346
Carew, M. E., 587
Carpintero, D. L., 694
Carstens, B. C., 165
Chaiwong, T., 113
Chattrukul, U., 887
Chen, B.-H., 475
Chi, H., 475
Chi, Y., 470
Choate, B. A., 496
Choi, W.-Y., 636
Cisneros, R., 960
Cognato, A. I., 931
Conn, J. E., 908
Cordo, H. A., 66
Coşkun, M., 336
Cui, L., 173

D

Dabert, J., 13
D'Adamo, P., 236
Danley, P. D., 219
Dastur, S. R., 989
Davis, A. K., 682
Delprat, A., 119
DeWalt, R. E., 941
Díaz, E., 960
Diez-Rodríguez, G. I , 409
Disney, H., 113
Dorado, J., 643
Dronnet, S., 395
Duyck, P. F., 402

E

Ehrnsberger, R., 13
Ellersieck, M., 787
Emre, I., 336
Engel, M. S., 73
Evans, W. G., 738

F

Fang, J., 351
Favret, C., 941
Fereres, A., 763
Ferris, I., 1
Ferro, D. N., 989
Filippi, L., 134, 143
Flakus, A., 108
Fletcher, J., 770
Fontal-Cazalla, F., 63
Forister, M. L., 295
Foster, P. G., 918
Franz, G., 119
Fuller, S., 126
Funk, D. J., 711

G

Galun, R., 302
Galvão, C., 861
Garofalo, R. S., 329
Garrido, A., 747
Gera, A., 351
Gillette, N., 931
Godoy, C., 444
Goodpasture, C., 227
Goolsby, J. H., 887
Goughnour, R. B., 703
Goulet, H., 565
Graves, W. R., 690, 926
Green, J., 309
Griswold, M. W., 673
Gut, L. J., 580

H

Habibi, J., 787
Hanski, I., 382
Harshman, D. B., 814
Hazel, W., 996
Heather, N., 1
Henderson, G., 340, 732
Hendrichs, J., 1
Hill, J. M., 552
Hironaka, M., 134, 143
Hirose, E., 515
Hirose, Y., 291
Hoagstrom, C. W., 241
Hoback, W. W., 973
Hoffmann, A. A., 587
Hogenhout, S. A., 820
Honda, J. Y., 291
Horton, D. R., 527, 608
Hoy, C. W., 756
Hussein, S., 227

I

Idris, A. M., 827
Infante, F., 483
Israely, N., 77, 302
Iwakuma, S., 143

J

Jacas, J.-A., 747
Jackson, B. C., 814
Jameson, M. L., 658
Johnson, D. G., 165
Jones, W. A., 322
Julien, M., 643
Jurberg, J., 861

K

Kankare, M., 382
Keena, M. A., 536

Kilpatrick, C. W., 908
Klag, J., 108
Knisley, C. B., 552, 973
Koç, H., 346
Kovarik, P. W., 245
Krecek, J., 273
Kritzman, A., 351
Kung, G.-A., 55
Kuyucu, N., 346

L

Lagos, D., 690
Laine, R. A., 340, 732
Langellotto, G. A., 205
Larivee, M., 951
Lasnier, J., 565
Lavi, U., 351
Lee, M.-M., 873
Lehr, M. A., 908
Lesage, L., 565
Lewis, M. L., 181
Lewis, T. M., 527, 608
Lewis, W. J., 595
Li, C., 918
Lin, H., 329
Liu, T.-X., 726
Liu, Y., 340, 732
Llácer, E., 747
Logarzo, G. A., 694
Lopes, J.R.S., 775
Lounibos, L. P., 673
Lourenço-de-Oliveira, R., 838
Lozada, M., 236
Lubkin, S. H., 73
Luis, A., 227

M

Madden, L. V., 756
Mahroof, R., 100
Maier, C. T., 214
Mallet, J., 873
Mao, L., 340, 732
Mariani, R., 66
Marino de Remes Lenicov, A. M., 66
Marsh, P. M., 636
Marshall, K., 996
Mathis, W. N., 431
Matthews, R. W., 595
McAloon, F. M., 861
McPheron, B. A., 173
McPherson, J. E., 18, 282
Messenger, M. T., 395, 980
Miller, J. R., 580
Miller, S. A., 756
Miller, T. A., 814
Mironov, S. V., 13
Mitchell, F., 770
Miura, K., 96, 470
Molina, J. M., 191
Moreno, A., 763
Morgante, J. S., 367
Morse, G. E., 629
Motta, M. A., 838

N

Nakamura, K., 96
Nault, L. R., 756
Nava, D. E., 409
Nieves-Aldrey, J. L., 853
Niyazi, N., 119
Nomakuchi, S., 134, 143
Normark, B. B., 629
Norrbom, A. L., 252
Novoa, F., 558, 896

O

Oku, K., 625
Oman, S. D., 77
O'Neil, R. J., 690, 926
Osakabe, M., 625
Ovruski, S. M., 252
Özalp, P., 336

P

Pair, S., 770
Palacios, I., 763
Panizzi, A. R., 515
Patterson, J., 861
Peck, S. B., 951
Pemberton, R., 887
Peredo, L. C., 458
Pérez, J., 483
Perondini, A.L.P., 367
Pettigrove, V., 587
Piangjai, S., 113
Polavarapu, S., 197, 966
Pollak, N., 351
Poore, M., 309
Porter, A. H., 989
Porter, S. D., 85
Postali Parra, J. R., 409
Provencher, L. M., 629
Pujade-Villar, J., 63
Purcell, A. H., 775

Q

Quilici, S., 402

R

Radtke, M. G., 548
Rakitov, R. A., 444
Rasnitsyn, A. P., 503
Reid, C.M.A., 243, 243
Reithel, J. S., 491
Reitz, S. R., 245
Riddick, E. W., 615
Rieske, L. K., 496
Rimbut, S., 887
Ringo, J., 601
Robinson, A. S., 119
Rocha, D. S., 861
Romero-Severson, J., 359
Rosenheim, J. A., 205
Ros-Farraré, P., 63
Rost, M. M., 108
Rota, J., 37
Rousse, P., 402
Ruberson, J. R., 595
Ryckewaert, P., 402

S

Sakamaki, Y., 470
Sallum, M.A.M., 918
Sanborn, A. F., 187
Sánchez, L., 960
Sánchez-Peña, S. R., 151
Santos, E. R., 119
Schaefer, C. W., 515, 861
Schaefer, P. W., 615
Scheffer, S. J., 181
Scheffrahn, R. H., 273, 395, 980
Scherer, A. M., 552
Schliserman, P., 252
Schwarz, M. P., 126
Schwarz, P. M., 126
Selivon, D., 367
Sétamou, M., 322
Shapiro, A. M., 227
Shaw, K. L., 219
Shelly, T., 755
Sheppard, W. S., 165
Shurtz, R. A., 18
Silva-Torres, C.S.A., 595
Simões Bento, J. M., 409
Skinner, K. F., 973
Solis, M. A., 887
Sorenson, C., 309
Sosa, A. J., 66, 643
Stelinski, L. L., 580
Stoffolano, J. G., Jr., 329
Stone, N., 996
Strazanac, J. S., 636
Stuart, J. J., 359
Subramanyam, B., 100
Sukontason, K., 113
Sukontason, K. L., 113
Suludere, Z., 346
Swoboda, K. A., 658
Szalanski, A. L., 273, 395, 980

T

Takafuji, A., 625
Talyn, B., 601
Taylor, L., 291
Teixeira, L.A.F., 197, 966
Thagong, A., 887
Tierney, S. M., 126
Tokuda, M., 259

U

Ueno, H., 96
Underwood, D.L.A., 227
Urbaneja, A., 747

V
van Achterberg, C., 636
van Nouhuys, S., 382
Vega, F. E., 359, 483
Vincent, C., 565
Voegtlin, D. J., 690, 926
Vogtsberger, R. C., 113

W
Walton, R. K., 682
Washburn, S., 309
Watkins, R. A., 242, 754
Watson, D. W., 309
Wayadande, A., 755, 770
Webb, D. W., 941
Weeks, A. R., 629
Whinnett, A., 873
Wilkerson, R. C., 908, 918
Will, K. W., 415
Williams, L., III, 694
Williams, M. R., 205
Williamson, G. B., 548
Willmott, K. R., 873
Winotai, A., 887
Wornoayporn, V., 119
Wright, T., 887
Wyatt, A., 996

Y
Yan, F., 787
Yang, Z.-Q., 636
Yano, S., 625
Yashiro, N., 291
Yee, W. L., 703
Yen, S.-H., 887
Yin, C.-M., 329
Yonash, N., 351
Yoo, H.J.S., 690, 926
Yu, J.-Z., 475
Yukawa, J., 259

Z
Zack, R. S., 165
Zahniser, J. N., 653
Zammit, J., 126
Zatwarnicki, T., 431
Zhu, K. Y., 100
Ziv, Y., 77, 302
Zmudzinska, A., 565
Zúñiga, G., 960

CUMULATIVE KEY WORD INDEX
Annals of Entomology
Volume 98, 2005

A

Acanthocephalini,
bamboo, 48
Cervantistellus, 48
Coreidae, 48
Mexico, 48
aculeus,
chromosomes, 367
Haldane's rule, 367
isozymes, 367
wings, 367
Adephaga,
Aspidytidae, 417
Dytiscoidea, 417
larval morphology, 417
phylogeny, 417
adult emergence,
Pimpla turionellae, 336
sex ratio, 336
vitamin E, 336
aggregations,
Harmonia axyridis, 615
infection, 615
overwinter, 615
Pennsylvania, 615
Agrilus planipennis,
biocontrol, 636
new species, 636
parasitoid, 636
Spathius agrili, 636
agromyzid,
cryptic species, 181
invasive species, 181
leafminer, 181
phylogeography, 181
Aleyrodidae,
cryptic species, 827
genetic variation, 827
species complex, 827
virus vector, 827
alligator weed,
Alternanthera philoxeroides, 643
Alticinae, 643
biological control, 643
Systena nitentula, 643
allochronic isolation,
Callophrys, 295
Cupressaceae, 295
host race, 295
Lycaenidae, 295
Alternanthera philoxeroides,
alligator weed, 643
Alticinae, 643
biological control, 643
Systena nitentula, 643
Alticinae,
alligator weed, 643
Alternanthera philoxeroides, 643
biological control, 643
Systena nitentula, 643
amplified fragment length polymorphism, 351, 359
biodiversity, 351
Broca, 359
DNA fingerprinting, 359
genetic relationship, 351
invasive species, 359
polymorphism, 351
pseudo-arrhenotoky, 359
Anastrepha,
C3C5 region, 173
CCID, 173
period, 173
Tephritidae, 173
Anopheles albitarsis Complex,
Culicidae, 918
molecular phylogeny, 918
Anopheles albitarsis s.l.,
cryptic species, 908
mtDNA *COI* gene, 908
RAPD-PCR, 908
Anoplophora glabripennis,
artificial diets, 536
iron, 536
rearing, 536
anoxia,
Cicindelidae, 973
flooding., 973
soil moisture, 973
tiger beetle, 973
antenna,
coeloconic sensilla, 113
maxillary palp, 113
Megaselia scalaris, 113
ultrastructure, 113
ant-tending,
mutualism, 491
nitrogen, 491
Aphis glycines, 690, 926
buckthorn, 690, 926
primary host plant, 926
Rhamnus, 690
Rhamnus alnifolia, 926
Rhamnus cathartica, 926
soybean aphid winter host, 690
Aphis gossypii,
Lemnia biplagiata, 475
Life table, 475
predation, 475
Aphodius,
cattle, 309
coastal-plain, 309
Onthophagus, 309
piedmont, 309
apple,
distribution, 214
emergence, 214
hosts, 214
seasonal flight period, 214
aquatic insect conservation,
Illinois, 941
Plecoptera, 941
arboreal termite,
distribution, 273
taxonomy, 273
Argentina,
biological control, 85
parasitoid, 85
Pseudacteon, 85
Solenopsis, 85
armored scale insects,
Aspidiotini, 629
endosymbiosis, 629
parthenogenesis, 629
thelytoky, 629
arthropods,
disinfestation, 1
irradiation, 1
sterile insect technique, 1
sterilization, 1
artificial diets,
Anoplophora glabripennis, 536
iron, 536
rearing, 536
Ascia monuste,
biology, 726
butterfly, 726
cruciferous vegetables, 726
Aspidiotini,
armored scale insects, 629
endosymbiosis, 629
parthenogenesis, 629
thelytoky, 629
Aspidytidae,
Adephaga, 417
Dytiscoidea, 417
larval morphology, 417
phylogeny, 417
aster leafhopper,
aster yellows phytoplasma, 756
beta-binomial distribution, 756
insect traps, 756
aster yellows phytoplasma,
aster leafhopper, 756
beta-binomial distribution, 756
insect traps, 756
Athysanini,
Deltocephalinae, 653
Luheriini, 653
new genus, 653
new species, 653
attractiveness,
pheromones, 236
social wasps, 236
yellowjackets, 236

B

B chromosomes,
biased sex ratio, 227
spermatogenesis, 227
bacteria,
biofilm, 775
Cicadellidae, 775
Hemiptera, 775
plant disease, 775
Bactrocera cucurbitae,
field cages, 402
methodology, 402
protein baits, 402
Tephritidae, 402
bamboo, 48, 444
Acanthocephalini, 48
brochosomes, 444
Cervantistellus, 48
Coreidae, 48

leafhoppers, 444
Mexico, 48
new species, 444
oviposition, 444
beta diversity,
biogeography, 558
Chrysomelidae, 558
Galicia, 558
inventory completeness, 558
beta-binomial distribution,
aster leafhopper, 756
aster yellows phytoplasma, 756
insect traps, 756
biased sex ratio,
B chromosomes, 227
spermatogenesis, 227
biocontrol,
Agrilus planipennis, 636
new species, 636
parasitoid, 636
Spathius agrili, 636
biodiversity, 351, 565, 931
amplified fragment length polymorphism, 351
Coleoptera, 931
conservation genetics, 931
Curculionoidea, 565
genetic relationship, 351
nested clade analysis, 931
phylogeography, 931
polymorphism, 351
vineyard, 565
weevil, 565
biofilm,
bacteria, 775
Cicadellidae, 775
Hemiptera, 775
plant disease, 775
biogeography, 503, 558, 951
beta diversity, 558
Chrysomelidae, 558
Florida, 951
fossil insects, 503
Galicia, 558
inventory completeness, 558
Notiothauma reedi, 503
Okanagan Highlands, 503
over-water dispersal, 951
Paleogene, 503
peninsula effect, 951
West Indies, 951
biological control, 66, 85, 643, 747
alligator weed, 643
Alternanthera philoxeroides, 643
Alticinae, 643
Argentina, 85
Eichhornia crassipes, 66
Eulophidae, 747
Galeopsomyia fausta, 747
immature stages, 66
larval development, 747
larval morphology, 747
Megamelus scutellaris, 66
parasitoid, 85
Pseudacteon, 85
Solenopsis, 85
Systena nitentula, 643
water hyacinth, 66
biology, 515, 726
Ascia monuste, 726
butterfly, 726
Cardiospermum halicacabum, 515
cruciferous vegetables, 726
Rhopalidae, 515
Sapindaceae, 515
setal patterns, 515
biomass,
length–mass relationship, 548
Neotropical Scarabaeidae, 548
volume, 548
volume–mass relationship, 548
biometric data,
developmental biology, 322
Homalodisca coagulata, 322
life table parameters, 322
sharpshooter, 322
bitter cherry,
black hawthorn, 703
cascara, 703
cherry laurel, 703
Rhagoletis indifferens, 703
black hawthorn,
bitter cherry, 703
cascara, 703
cherry laurel, 703
Rhagoletis indifferens, 703
blueberry maggot, 197, 966
20-hydroxyecdysone, 197
diapause, 197, 966
heat stress, 197, 966
Hsp70, 966
Rhagoletis mendax, 966
body size,
fecundity, 191
oviposition, 191
pupal weight, 191
reproduction, 191
bottom-up,
coexistence, 673
mosquitoes, 673
top-down, 673
treehole, 673
Brevicoryne brassicae,
CaMV, 763
inoculation, 763
Myzus persicae, 763
vector transmission, 763
Broca,
amplified fragment length polymorphism, 359
DNA fingerprinting, 359
invasive species, 359
pseudo-arrhenotoky, 359
brochosomes,
bamboo, 444
leafhoppers, 444
new species, 444
oviposition, 444
Brochymena,
male genitalia, 282
Parabrochymena, 282
buckthorn, 690, 926
Aphis glycines, 690, 926
primary host plant, 926
Rhamnus, 690
Rhamnus alnifolia, 926
Rhamnus cathartica, 926
soybean aphid winter host, 690
butterfly,
Ascia monuste, 726
biology, 726
cruciferous vegetables, 726

C

^{13}C,
^{15}N, 205
parasitoids, 205
Rhopalomyia californica, 205
stable isotopes, 205
C3C5 region,
Anastrepha, 173
CCID, 173
period, 173
Tephritidae, 173
Callophrys,
allochronic isolation, 295
Cupressaceae, 295
host race, 295
Lycaenidae, 295
Campephilus principalis,
feather mites, 13
new species, 13
Pterotrogus, 13
Camptosomata,
Chlamisini, 711
insect architecture, 711
insect herbivores, 711
oviposition, 711
CaMV,
Brevicoryne brassicae, 763
inoculation, 763
Myzus persicae, 763
vector transmission, 763
Cardiospermum halicacabum,
biology, 515
Rhopalidae, 515
Sapindaceae, 515
setal patterns, 515
cascara,
bitter cherry, 703
black hawthorn, 703
cherry laurel, 703
Rhagoletis indifferens, 703
caste differentiation,
Formosan subterranean termite, 732
JH titer, 732
proximate mechanism, 732
soldier proportion, 732
caste regulation,
Coptotermes formosanus, 340
juvenile hormone, 340
soldier formation, 340
cattle,
Aphodius, 309
coastal-plain, 309
Onthophagus, 309
piedmont, 309
CCID,
Anastrepha, 173
C3C5 region, 173
period, 173
Tephritidae, 173
cell culture,
Dendroctonus, 960
larvae, 960
midgut, 960
Celtis,
host plants, 252
Rhagoletotrypeta cubensis n. sp., 252
taxonomy, 252
Tephritidae, 252

Ceratitis capitata, 77, 302
invasion, 77
large-scale population dynamics, 302
local extinction, 302
long-distance dispersal, 302
overwintering, 77
population distribution pattern, 77
recolonization, 302
spatial patterns, 77
Cercantistellus,
Acanthocephalini, 48
bamboo, 48
Coreidae, 48
Mexico, 48
Cetoniinae,
Dasycalgus, 658
Scarabaeidae, 658
Valgini, 658
Valgus, 658
Chagas disease,
Linshcosteus karupus, 861
morphology, 861
systematics, 861
Triatominae, 861
chemoreception,
flesh fly, 595
host location, 595
leafcutting bee, 595
parasitism, 595
cherry laurel,
bitter cherry, 703
black hawthorn, 703
cascara, 703
Rhagoletis indifferens, 703
Chironomidae,
Cladopelma, 587
DNA markers, 587
mitochondrial DNA, 587
taxonomy, 587
Chlamisini,
Camptosomata, 711
insect architecture, 711
insect herbivores, 711
oviposition, 711
chorion,
morphology, 346
SEM, 346
terminal filament, 346
Tipula, 346
chromosomes,
aculeus, 367
Haldane's rule, 367
isozymes, 367
wings, 367
Chrysomelidae, 558, 896
beta diversity, 558
biogeography, 558
female genitalia, 896
Galicia, 558
inventory completeness, 558
Neocrepidodera ferruginea group, 896
Neocrepidodera impressa group, 896
new species, 896
cicada,
Costa Rica, 187
Fidicina, 187
new species, 187
taxonomy, 187
Cicadellidae, 165, 775
bacteria, 775
biofilm, 775
Errhomus, 165
Erronus, 165
Hemiptera, 775
leafhopper, 165
molecular systematics, 165
plant disease, 775
Cicindelidae, 552, 973
anoxia, 973
flooding., 973
recovery, 552
soil moisture, 973
tiger beetle, 552, 973
translocation, 552
citrus root weevil,
Oestrophasia (*Cenosoma*) *sabroskyi*, 245
parasitoid, 245
Tachinidae, 245
Cladopelma,
Chironomidae, 587
DNA markers, 587
mitochondrial DNA, 587
taxonomy, 587
coastal-plain,
Aphodius, 309
cattle, 309
Onthophagus, 309
piedmont, 309
coeloconic sensilla,
antenna, 113
maxillary palp, 113
Megaselia scalaris, 113
ultrastructure, 113
coexistence,
bottom-up, 673
mosquitoes, 673
top-down, 673
treehole, 673
coffee berry borer,
fungi, 483
Hypothenemus, 483
mutualism, 483
Coleoptera, 73, 931
biodiversity, 931
conservation genetics, 931
fossil insects, 73
nested clade analysis, 931
Permian, 73
phylogeography, 931
Wellington Formation, 73
competition,
kleptoparasitism, 134
parental care, 134
provisioning, 134
risk-sensitive behavior, 134
conditional strategy,
polyphenism, 996
pupal color, 996
pupation sites, 996
conservation genetics,
biodiversity, 931
Coleoptera, 931
nested clade analysis, 931
phylogeography, 931
Coptotermes formosanus,
caste regulation, 340
juvenile hormone, 340
soldier formation, 340
Coreidae,
Acanthocephalini, 48
bamboo, 48
Cervantistellus, 48
Mexico, 48
corn stunt spiroplasma,
Dalbulus maidis, 820
immunofluorescence confocal laser scanning microscopy, 820
mollicutes, 820
Spiroplasma kunkelii, 820
Cornus controversa,
Elaeocarpus sylvestris, 259
gall midge, 259
Pseudasphondylia elaeocarpi, 259
Pseudasphondylia kiritanii, 259
Costa Rica,
cicada, 187
Fidicina, 187
new species, 187
taxonomy, 187
cruciferous vegetables,
Ascia monuste, 726
biology, 726
butterfly, 726
cryptic species, 181, 382, 608, 827, 908
agromyzid, 181
Aleyrodidae, 827
Anopheles albitarsis s.l., 908
female resistance, 608
genetic variation, 827
geographic variation, 608
Insecta, 608
insemination, 608
invasive species, 181
leafminer, 181
Melitaeini, 382
microsatellites, 382
mtDNA, 382
mtDNA *COI* gene, 908
parasitoids, 382
phylogeography, 181
RAPD-PCR, 908
species complex, 827
virus vector, 827
Culicidae, 838, 918
Anopheles albitarsis Complex, 918
molecular phylogeny, 918
new species, 838
Sabethini, 838
systematics, 838
Wyeomyia, 838
Cupressaceae,
allochronic isolation, 295
Callophrys, 295
host race, 295
Lycaenidae, 295
Curculionoidea,
biodiversity, 565
vineyard, 565
weevil, 565
Cydia pomonella,
electroantennogram, 580
Grapholita molesta, 580
long-lasting adaptation, 580
Pandemis pyrusana, 580
Cynipidae,
Iraella, 853
new species, 853
Papaver, 853
Spain, 853

D

Dalbulus maidis,
corn stunt spiroplasma, 820
immunofluorescence confocal laser scanning microscopy, 820
mollicutes, 820
Spiroplasma kunkelii, 820
Danaus plexippus,
endangered biological phenomenon, 682
North America, 682
population census, 682
seasonal migration, 682
Dasyvalgus,
Cetoniinae, 658
Scarabaeidae, 658
Valgini, 658
Valgus, 658
Deltocephalinae,
Athysanini, 653
Luheriini, 653
new genus, 653
new species, 653
Dendroctonus,
cell culture, 960
larvae, 960
midgut, 960
descriptions,
laboratory rearing, 18
life history, 18
nymphs, 18
Sinea spinipes, 18
development,
ecdysis, 219
Laupala, 219
stridulation, 219
developmental biology,
biometric data, 322
Homalodisca coagulata, 322
life table parameters, 322
sharpshooter, 322
diapause, 197, 575, 966
20-hydroxyecdysone, 197
blueberry maggot, 197, 966
Dolycoris baccarum, 575
fatty acid composition, 575
heat stress, 197, 966
Hsp70, 966
Piezodorus lituratus, 575
prediapause, 575
Rhagoletis mendax, 966
Diptera,
Discomyza, 431
Ephydridae, 431
malacophagy, 431
shore flies, 431
Discomyza,
Diptera, 431
Ephydridae, 431
malacophagy, 431
shore flies, 431
disinfestation,
arthropods, 1
irradiation, 1
sterile insect technique, 1
sterilization, 1
dispersal capacity,
mating status, 625
mean crowding, 625
Tetranychus kanzawai, 625
trail, 625
distribution, 214, 273
apple, 214
arboreal termite, 273
emergence, 214
hosts, 214
seasonal flight period, 214
taxonomy, 273
DNA fingerprinting,
amplified fragment length polymorphism, 359
Broca, 359
invasive species, 359
pseudo-arrhenotoky, 359
DNA markers,
Chironomidae, 587
Cladopelma, 587
mitochondrial DNA, 587
taxonomy, 587
DNA sequence, 395, 980
16S, 980
genetic variation, 395, 980
Reticulitermes, 395, 980
termite, 395, 980
Dohrniphora,
Neotropical, 55
Phoridae, 55
taxonomy, 55
Dolycoris baccarum,
diapause, 575
fatty acid composition, 575
Piezodorus lituratus, 575
prediapause, 575
dorsal setae on A9,
La Selva Biological Station, 37
Moraceae, 37
pupal dorsal spines, 37
Tortricidae, 37
Drosophila melanogaster,
female receptivity, 601
ovaries, 601
oviposition, 601
precocene, 601
Dytiscoidea,
Adephaga, 417
Aspidytidae, 417
larval morphology, 417
phylogeny, 417

E

eastern tent caterpillar,
life history, 496
mare reproductive loss syndrome, 496
mortality, 496
ecdysis,
development, 219
Laupala, 219
stridulation, 219
egg parasitoid,
Hymenoptera, 291
Insecta, 291
taxonomy, 291
Trichogrammatidae, 291
Eichhornia crassipes,
biological control, 66
immature stages, 66
Megamelus scutellaris, 66
water hyacinth, 66
Elaeocarpus sylvestris,
Cornus controversa, 259
gall midge, 259
Pseudasphondylia elaeocarpi, 259
Pseudasphondylia kiritanii, 259
electrical penetration graph,
electronic monitoring of insect feeding, 787
Homalodisca coagulata, 787
Pierce's disease, 787
probing, 787
electroantennogram,
Cydia pomonella, 580
Grapholita molesta, 580
long-lasting adaptation, 580
Pandemis pyrusana, 580
electronic monitoring of insect feeding,
electrical penetration graph, 787
Homalodisca coagulata, 787
Pierce's disease, 787
probing, 787
electrophoretic analysis,
F1 sterility, 470
hybridization, 470
L. trifolii, 470
Liriomyza sativae, 470
embryogenesis,
gonads, 108
PGCs, 108
emergence,
apple, 214
distribution, 214
hosts, 214
seasonal flight period, 214
endangered biological phenomenon,
Danaus plexippus, 682
North America, 682
population census, 682
seasonal migration, 682
endosymbiosis,
armored scale insects, 629
Aspidiotini, 629
parthenogenesis, 629
thelytoky, 629
Ephydridae,
Diptera, 431
Discomyza, 431
malacophagy, 431
shore flies, 431
Errhomus,
Cicadellidae, 165
Erronus, 165
leafhopper, 165
molecular systematics, 165
Erronus,
Cicadellidae, 165
Errhomus, 165
leafhopper, 165
molecular systematics, 165
Eucoilinae,
Figitidae, 63
Hymenoptera, 63
Perischus, 63
Eulophidae,
biological control, 747
Galeopsomyia fausta, 747
larval development, 747
larval morphology, 747
eusociality,
Madagascar, 126
mass provisioning, 126
phylogenetics, 126
sociality, 126

F

F1 sterility,
electrophoretic analysis, 470
hybridization, 470
L. trifolii, 470
Liriomyza sativae, 470
fatty acid composition,
diapause, 575
Dolycoris baccarum, 575
Piezodorus lituratus, 575
prediapause, 575
feather mites,
Campephilus principalis, 13
new species, 13
Pterotrogus, 13
fecundity,
body size, 191
oviposition, 191
pupal weight, 191
reproduction, 191
feeding behavior,
plant disease, 814
vector, 814
female genitalia,
Chrysomelidae, 896
Neocrepidodera ferruginea group, 896
Neocrepidodera impressa group, 896
new species, 896
female receptivity,
Drosophila melanogaster, 601
ovaries, 601
oviposition, 601
precocene, 601
female resistance,
cryptic species, 608
geographic variation, 608
Insecta, 608
insemination, 608
Fidicina,
cicada, 187
Costa Rica, 187
new species, 187
taxonomy, 187
field cage, 119, 402
Bactrocera cucurbitae, 402
genetic marker, 119
Mediterranean fruit fly, 119
methodology, 402
protein baits, 402
Tephritidae, 402
Figitidae,
Eucoilinae, 63
Hymenoptera, 63
Perischus, 63
flesh fly,
chemoreception, 595
host location, 595
leafcutting bee, 595
parasitism, 595
flooding,
anoxia, 973
Cicindelidae, 973
soil moisture, 973
tiger beetle, 973
Florida,
biogeography, 951
over-water dispersal, 951
peninsula effect, 951
West Indies, 951
Formosan subterranean termite,
caste differentiation, 732
JH titer, 732
proximate mechanism, 732
soldier proportion, 732
fossil insects, 73, 503
biogeography, 503
Coleoptera, 73
Notiothauma reedi, 503
Okanagan Highlands, 503
Paleogene, 503
Permian, 73
Wellington Formation, 73
fungi,
coffee berry borer, 483
Hypothenemus, 483
mutualism, 483
Fungus-growing ants,
mycophagy, 151
origin, 151
symbiosis, 151

G

Galeopsomyia fausta,
biological control, 747
Eulophidae, 747
larval development, 747
larval morphology, 747
Galicia,
beta diversity, 558
biogeography, 558
Chrysomelidae, 558
inventory completeness, 558
gall midge,
Cornus controversa, 259
Elaeocarpus sylvestris, 259
Pseudasphondylia elaeocarpi, 259
Pseudasphondylia kirʻtanii, 259
genetic marker,
field cage, 119
Mediterranean fruit fly, 119
genetic relationship,
amplified fragment length polymorphism, 351
biodiversity, 351
polymorphism, 351
genetic variation, 395, 827, 980
16S, 980
Aleyrodidae, 827
cryptic species, 827
DNA sequence, 395, 980
Reticulitermes, 395, 980
species complex, 827
termite, 395, 980
virus vector, 827
genitalia,
Insecta, 527
mating isolation, 527
morphometrics, 527
sexual selection, 527
geographic distribution,
Miridae, 694
plant bugs host plants, 694
species richness, 694
geographic variation,
cryptic species, 608
female resistance, 608
Insecta, 608
insemination, 608
gonads,
embryogenesis, 108
PGCs, 108
Grapholita molesta,
Cydia pomonella, 580
electroantennogram, 580
long-lasting adaptation, 580
Pandemis pyrusana, 580

H

20-hydroxyecdysone,
blueberry maggot, 197
diapause, 197
heat stress, 197
Haldane's rule,
aculeus, 367
chromosomes, 367
isozymes, 367
wings, 367
Harmonia axyridis, 96, 615
aggregations, 615
infection, 615
ladybird beetle, 96
male-killing, 96
overwinter, 615
Pennsylvania, 615
prevalence, 96
Spiroplasma, 96
heat shock proteins,
heat treatment, 100
thermotolerance, 100
Tribolium castaneum, 100
heat stress, 197, 966
20-hydroxyecdysone, 197
blueberry maggot, 197, 966
diapause, 197, 966
Hsp70, 966
Rhagoletis mendax, 966
heat treatment,
heat shock proteins, 100
thermotolerance, 100
Tribolium castaneum, 100
Hemiptera,
bacteria, 775
biofilm, 775
Cicadellidae, 775
plant disease, 775
Heteroptera, 458, 770
Mexico, 458
Myodocha, 458
propagative bacteria, 770
Rhyparochromidae, 458
squash bug, 770
transmission, 770
Homalodisca coagulata, 322, 787
biometric data, 322
developmental biology, 322
electrical penetration graph, 787
electronic monitoring of insect feeding, 787
life table parameters, 322
Pierce's disease, 787
probing, 787
sharpshooter, 322
host,
apple, 214
distribution, 214
emergence, 214
seasonal flight period, 214
host location,
chemoreception, 595
flesh fly, 595
leafcutting bee, 595
parasitism, 595

host plants,
Celtis, 252
Rhagoletotrypeta cubensis n. sp., 252
taxonomy, 252
Tephritidae, 252
host race,
allochronic isolation, 295
Callophrys, 295
Cupressaceae, 295
Lycaenidae, 295
Hsp70,
blueberry maggot, 966
diapause, 966
heat stress, 966
Rhagoletis mendax, 966
hybridization,
electrophoretic analysis, 470
F1 sterility, 470
L. trifolii, 470
Liriomyza sativae, 470
Hymenoptera, 63, 291
egg parasitoid, 291
Eucoilinae, 63
Figitidae, 63
Insecta, 291
Perischus, 63
taxonomy, 291
Trichogrammatidae, 291
Hypothenemus,
coffee berry borer, 483
fungi, 483
mutualism, 483

I

Illinois,
aquatic insect conservation, 941
Plecoptera, 941
immature stages,
biological control, 66
Eichhornia crassipes, 66
Megamelus scutellaris, 66
water hyacinth, 66
immunofluorescence confocal laser scanning microscopy,
corn stunt spiroplasma, 820
Dalbulus maidis, 820
mollicutes, 820
Spiroplasma kunkelii, 820
infection,
aggregations, 615
Harmonia axyridis, 615
overwinter, 615
Pennsylvania, 615
infrared radiation receptor,
Melanophila, 738
phasic receptor, 738
sensor structure, 738
transduction, 738
inoculation,
Brevicoryne brassicae, 763
CaMV, 763
Myzus persicae, 763
vector transmission, 763
insect architecture,
Camptosomata, 711
Chlamisini, 711
insect herbivores, 711
oviposition, 711
insect herbivores,
Camptosomata, 711
Chlamisini, 711
insect architecture, 711
oviposition, 711
insect traps,
aster leafhopper, 756
aster yellows phytoplasma, 756
beta-binomial distribution, 756
Insecta, 291, 527, 608
cryptic species, 608
egg parasitoid, 291
female resistance, 608
genitalia, 527
geographic variation, 608
Hymenoptera, 291
insemination, 608
mating isolation, 527
morphometrics, 527
sexual selection, 527
taxonomy, 291
Trichogrammatidae, 291
insecticide resistance management,
Leptinotarsa decemlineata, 989
mating behavior, 989
sperm precedence, 989
insemination,
cryptic species, 608
female resistance, 608
geographic variation, 608
Insecta, 608
invasion,
Ceratitis capitata, 77
overwintering, 77
population distribution pattern, 77
spatial patterns, 77
invasive species, 181, 359
agromyzid, 181
amplified fragment length polymorphism, 359
Broca, 359
cryptic species, 181
DNA fingerprinting, 359
leafminer, 181
phylogeography, 181
pseudo-arrhenotoky, 359
inventory completeness,
beta diversity, 558
biogeography, 558
Chrysomelidae, 558
Galicia, 558
Iraella,
Cynipidae, 853
new species, 853
Papaver, 853
Spain, 853
iron,
Anoplophora glabripennis, 536
artificial diets, 536
rearing, 536
irradiation,
arthropods, 1
disinfestation, 1
sterile insect technique, 1
sterilization, 1
isozymes,
aculeus, 367
chromosomes, 367
Haldane's rule, 367
wings, 367
Ithomiinae,
Lepidoptera, 873
molecular phylogenetics, 873
Tektin, 873

J

JH titer,
caste differentiation, 732
Formosan subterranean termite, 732
proximate mechanism, 732
soldier proportion, 732
juvenile hormone,
caste regulation, 340
Coptotermes formosanus, 340
soldier formation, 340

K

kleptoparasitism,
competition, 134
parental care, 134
provisioning, 134
risk-sensitive behavior, 134

L

L. trifolii,
electrophoretic analysis, 470
F1 sterility, 470
hybridization, 470
Liriomyza sativae, 470
La Selva Biological Station,
dorsal setae on A9, 37
Moraceae, 37
pupal dorsal spines, 37
Tortricidae, 37
laboratory rearing,
descriptions, 18
life history, 18
nymphs, 18
Sinea spinipes, 18
ladybird beetle,
Harmonia axyridis, 96
male-killing, 96
prevalence, 96
Spiroplasma, 96
large-scale population dynamics,
Ceratitis capitata, 302
local extinction, 302
long-distance dispersal, 302
recolonization, 302
larvae,
cell culture, 960
Dendroctonus, 960
midgut, 960
larval development,
biological control, 747
Eulophidae, 747
Galeopsomyia fausta, 747
larval morphology, 747
larval morphology, 417, 747
Adephaga, 417
Aspidytidae, 417
biological control, 747
Dytiscoidea, 417
Eulophidae, 747
Galeopsomyia fausta, 747
larval development, 747
phylogeny, 417
lateral neurosecretory cells,
liver-fed flies, 329
median neurosecretory cells, 329
midgut hormone, 329
sugar-fed flies, 329
Laupala,
development, 219
ecdysis, 219
stridulation, 219

leafcutting bee,
chemoreception, 595
flesh fly, 595
host location, 595
parasitism, 595
leafhopper, 165, 444
bamboo, 444
brochosomes, 444
Cicadellidae, 165
Errhomus, 165
Erronus, 165
molecular systematics, 165
new species, 444
oviposition, 444
leafminer,
agromyzid, 181
cryptic species, 181
invasive species, 181
phylogeography, 181
Lemnia biplagiata,
Aphis gossypii, 475
Life table, 475
predation, 475
length–mass relationship,
biomass, 548
Neotropical Scarabaeidae, 548
volume, 548
volume–mass relationship, 548
Lepidoptera,
Ithomiinae, 873
molecular phylogenetics, 873
Tektin, 873
Lepidoptera oviposition,
Persea americana, 409
Leptinotarsa decemlineata,
insecticide resistance management, 989
mating behavior, 989
sperm precedence, 989
life history, 18, 496
descriptions, 18
eastern tent caterpillar, 496
laboratory rearing, 18
mare reproductive loss syndrome, 496
mortality, 496
nymphs, 18
Sinea spinipes, 18
Life table,
Aphis gossypii, 475
Lemnia biplagiata, 475
predation, 475
life table parameters,
biometric data, 322
developmental biology, 322
Homalodisca coagulata, 322
sharpshooter, 322
Linshcosteus karupus,
Chagas disease, 861
morphology, 861
systematics, 861
Triatominae, 861
Liriomyza sativae,
electrophoretic analysis, 470
F1 sterility, 470
hybridization, 470
L. trifolii, 470
liver-fed flies,
lateral neurosecretory cells, 329
median neurosecretory cells, 329
midgut hormone, 329
sugar-fed flies, 329
local extinction,
Ceratitis capitata, 302
large-scale population dynamics, 302
long-distance dispersal, 302
recolonization, 302
long-distance dispersal,
Ceratitis capitata, 302
large-scale population dynamics, 302
local extinction, 302
recolonization, 302
long-lasting adaptation,
Cydia pomonella, 580
electroantennogram, 580
Grapholita molesta, 580
Pandemis pyrusana, 580
Luheriini,
Athysanini, 653
Deltocephalinae, 653
new genus, 653
new species, 653
Lycaenidae,
allochronic isolation, 295
Callophrys, 295
Cupressaceae, 295
host race, 295

M

Madagascar,
eusociality, 126
mass provisioning, 126
phylogenetics, 126
sociality, 126
malacophagy,
Diptera, 431
Discomyza, 431
Ephydridae, 431
shore flies, 431
male genitalia,
Brochymena, 282
Parabrochymena, 282
male-killing,
Harmonia axyridis, 96
ladybird beetle, 96
prevalence, 96
Spiroplasma, 96
mare reproductive loss syndrome,
eastern tent caterpillar, 496
life history, 496
mortality, 496
mass provisioning,
eusociality, 126
Madagascar, 126
phylogenetics, 126
sociality, 126
mating behavior,
insecticide resistance management, 989
Leptinotarsa decemlineata, 989
sperm precedence, 989
mating isolation,
genitalia, 527
Insecta, 527
morphometrics, 527
sexual selection, 527
mating status,
dispersal capacity, 625
mean crowding, 625
Tetranychus kanzawai, 625
trail, 625
maxillary palp,
antenna, 113
coeloconic sensilla, 113
Megaselia scalaris, 113
ultrastructure, 113
mean crowding,
dispersal capacity, 625
mating status, 625
Tetranychus kanzawai, 625
trail, 625
median neurosecretory cells,
lateral neurosecretory cells, 329
liver-fed flies, 329
midgut hormone, 329
sugar-fed flies, 329
Mediterranean fruit fly,
field cage, 119
genetic marker, 119
Megamelus scutellaris,
biological control, 66
Eichhornia crassipes, 66
immature stages, 66
water hyacinth, 66
Megaselia scalaris,
antenna, 113
coeloconic sensilla, 113
maxillary palp, 113
ultrastructure, 113
Melanophila,
infrared radiation receptor, 738
phasic receptor, 738
sensor structure, 738
transduction, 738
Melitaeini,
cryptic species, 382
microsatellites, 382
mtDNA, 382
parasitoids, 382
methodology,
Bactrocera cucurbitae, 402
field cages, 402
protein baits, 402
Tephritidae, 402
Mexico, 48, 458
Acanthocephalini, 48
bamboo, 48
Cervantistellus, 48
Coreidae, 48
Heteroptera, 458
Myodocha, 458
Rhyparochromidae, 458
microsatellites,
cryptic species, 382
Melitaeini, 382
mtDNA, 382
parasitoids, 382
midgut,
cell culture, 960
Dendroctonus, 960
larvae, 960
midgut hormone,
lateral neurosecretory cells, 329
liver-fed flies, 329
median neurosecretory cells, 329
sugar-fed flies, 329
Miridae,
geographic distribution, 694
plant bugs host plants, 694
species richness, 694
mitochondrial DNA,
Chironomidae, 587
Cladopelma, 587

DNA markers, 587
taxonomy, 587
molecular phylogenetics,
Ithomiinae, 873
Lepidoptera, 873
Tektin, 873
molecular phylogeny,
Anopheles albitarsis Complex, 918
Culicidae, 918
molecular systematics,
Cicadellidae, 165
Errhomus, 165
Erronus, 165
leafhopper, 165
mollicutes,
corn stunt spiroplasma, 820
Dalbulus maidis, 820
immunofluorescence confocal laser scanning microscopy, 820
Spiroplasma kunkelii, 820
Moraceae,
dorsal setae on A9, 37
La Selva Biological Station, 37
pupal dorsal spines, 37
Tortricidae, 37
morphology, 346, 861
Chagas disease, 861
chorion, 346
Linshcosteus karupus, 861
SEM, 346
systematics, 861
terminal filament, 346
Tipula, 346
Triatominae, 861
morphometrics,
genitalia, 527
Insecta, 527
mating isolation, 527
sexual selection, 527
mortality,
eastern tent caterpillar, 496
life history, 496
mare reproductive loss syndrome, 496
mosquitoes,
bottom-up, 673
coexistence, 673
top-down, 673
treehole, 673
mtDNA,
cryptic species, 382
Melitaeini, 382
microsatellites, 382
parasitoids, 382
mtDNA *COI* gene,
Anopheles albitarsis s.l., 908
cryptic species, 908
RAPD-PCR, 908
mutualism, 483, 491
ant-tending, 491
coffee berry borer, 483
fungi, 483
Hypothenemus, 483
nitrogen, 491
mycophagy,
Fungus-growing ants, 151
origin, 151
symbiosis, 151
Myodocha,
Heteroptera, 458
Mexico, 458
Rhyparochromidae, 458
Myzus persicae,
Brevicoryne brassicae, 763
CaMV, 763
inoculation, 763
vector transmission, 763

N

^{15}N,
^{13}C, 205
parasitoids, 205
Rhopalomyia californica, 205
stable isotopes, 205
Neocrepidodera ferruginea group,
Chrysomelidae, 896
female genitalia, 896
Neocrepidodera impressa group, 896
new species, 896
Neocrepidodera impressa group,
Chrysomelidae, 896
female genitalia, 896
Neocrepidodera ferruginea group, 896
new species, 896
Neotropical,
Dohrniphora, 55
Phoridae, 55
taxonomy, 55
Neotropical Scarabaeidae,
biomass, 548
length–mass relationship, 548
volume, 548
volume–mass relationship, 548
nested clade analysis,
biodiversity, 931
Coleoptera, 931
conservation genetics, 931
phylogeography, 931
new genus,
Athysanini, 653
Deltocephalinae, 653
Luheriini, 653
new species, 653
new species, 13, 187, 444, 636, 653, 838, 853, 896
Agrilus planipennis, 636
Athysanini, 653
bamboo, 444
biocontrol, 636
brochosomes, 444
Campephilus principalis, 13
Chrysomelidae, 896
cicada, 187
Costa Rica, 187
Culicidae, 838
Cynipidae, 853
Deltocephalinae, 653
feather mites, 13
female genitalia, 896
Fidicina, 187
Iraella, 853
leafhoppers, 444
Luheriini, 653
Neocrepidodera ferruginea group, 896
Neocrepidodera impressa group, 896
new genus, 653
oviposition, 444
Papaver, 853
parasitoid, 636
Pterotrogus, 13
Sabethini, 838
Spain, 853
Spathius agrili, 636
systematics, 838
taxonomy, 187
Wyeomyia, 838
nitrogen,
ant-tending, 491
mutualism, 491
North America,
Danaus plexippus, 682
endangered biological phenomenon, 682
population census, 682
seasonal migration, 682
Notiothauma reedi,
biogeography, 503
fossil insects, 503
Okanagan Highlands, 503
Paleogene, 503
nymphal independence,
parental care, 143
progressive provisioning, 143
shield bug, 143
nymphs,
descriptions, 18
laboratory rearing, 18
life history, 18
Sinea spinipes, 18

O

Oestrophasia (*Cenosoma*) *sabroskyi*,
citrus root weevil, 245
parasitoid, 245
Tachinidae, 245
Okanagan Highlands,
biogeography, 503
fossil insects, 503
Notiothauma reedi, 503
Paleogene, 503
Old World climbing fern,
pteridophagy, 887
Pyraloidea, 887
spiders, 887
tenebrionid beetles, 887
Onthophagus,
Aphodius, 309
cattle, 309
coastal-plain, 309
piedmont, 309
origin,
Fungus-growing ants, 151
mycophagy, 151
symbiosis, 151
ovaries,
Drosophila melanogaster, 601
female receptivity, 601
oviposition, 601
precocene, 601
over-water dispersal,
biogeography, 951
Florida, 951
peninsula effect, 951
West Indies, 951
overwinter, 77, 615
aggregations, 615
Ceratitis capitata, 77
Harmonia axyridis, 615
infection, 615
invasion, 77
Pennsylvania, 615

population distribution pattern, 77
spatial patterns, 77
oviposition, 191, 444, 601, 711
bamboo, 444
body size, 191
brochosomes, 444
Camptosomata, 711
Chlamisini, 711
Drosophila melanogaster, 601
fecundity, 191
female receptivity, 601
insect architecture, 711
insect herbivores, 711
leafhoppers, 444
new species, 444
ovaries, 601
precocene, 601
pupal weight, 191
reproduction, 191

P

Paleogene,
biogeography, 503
fossil insects, 503
Notiothauma reedi, 503
Okanagan Highlands, 503
Pandemis pyrusana,
Cydia pomonella, 580
electroantennogram, 580
Grapholita molesta, 580
long-lasting adaptation, 580
Papaver,
Cynipidae, 853
Iraella, 853
new species, 853
Spain, 853
Parabrochymena,
Brochymena, 282
male genitalia, 282
parasitism,
chemoreception, 595
flesh fly, 595
host location, 595
leafcutting bee, 595
parasitoid, 85, 205, 245, 382, 636
Agrilus planipennis, 636
Argentina, 85
biocontrol, 636
biological control, 85
^{13}C, 205
citrus root weevil, 245
cryptic species, 382
Melitaeini, 382
microsatellites, 382
mtDNA, 382
^{15}N, 205
new species, 636
Oestrophasia (*Cenosoma*) *sabroskyi*, 245
Pseudacteon, 85
Rhopalomyia californica, 205
Solenopsis, 85
Spathius agrili, 636
stable isotopes, 205
Tachinidae, 245
parental care, 134, 143
competition, 134
kleptoparasitism, 134
nymphal independence, 143
progressive provisioning, 143
provisioning, 134
risk-sensitive behavior, 134
shield bug, 143
parthenogenesis,
armored scale insects, 629
Aspidiotini, 629
endosymbiosis, 629
thelytoky, 629
peninsula effect,
biogeography, 951
Florida, 951
over-water dispersal, 951
West Indies, 951
Pennsylvania,
aggregations, 615
Harmonia axyridis, 615
infection, 615
overwinter, 615
period,
Anastrepha, 173
C3C5 region, 173
CCID, 173
Tephritidae, 173
Perischus,
Eucoilinae, 63
Figitidae, 63
Hymenoptera, 63
Permian,
Coleoptera, 73
fossil insects, 73
Wellington Formation, 73
Persea americana,
Lepidoptera oviposition, 409
PGCs,
embryogenesis, 108
gonads, 108
phasic receptor,
infrared radiation receptor, 738
Melanophila, 738
sensor structure, 738
transduction, 738
pheromones,
attractiveness, 236
social wasps, 236
yellowjackets, 236
Phoridae,
Dohrniphora, 55
Neotropical, 55
taxonomy, 55
phylogenetics,
eusociality, 126
Madagascar, 126
mass provisioning, 126
sociality, 126
phylogeny,
Adephaga, 417
Aspidytidae, 417
Dytiscoidea, 417
larval morphology, 417
phylogeography, 181, 931
agromyzid, 181
biodiversity, 931
Coleoptera, 931
conservation genetics, 931
cryptic species, 181
invasive species, 181
leafminer, 181
nested clade analysis, 931
piedmont,
Aphodius, 309
cattle, 309
coastal-plain, 309
Onthophagus, 309
Pierce's disease,
electrical penetration graph, 787
electronic monitoring of insect feeding, 787
Homalodisca coagulata, 787
probing, 787
Piezodorus lituratus,
diapause, 575
Dolycoris baccarum, 575
fatty acid composition, 575
prediapause, 575
Pimpla turionellae,
adult emergence, 336
sex ratio, 336
vitamin E, 336
plant bugs host plants,
geographic distribution, 694
Miridae, 694
species richness, 694
plant disease, 775, 814
bacteria, 775
biofilm, 775
Cicadellidae, 775
feeding behavior, 814
Hemiptera, 775
vector, 814
Plecoptera,
aquatic insect conservation, 941
Illinois, 941
polymorphism,
amplified fragment length polymorphism, 351
biodiversity, 351
genetic relationship, 351
polyphenism,
conditional strategy, 996
pupal color, 996
pupation sites, 996
population census,
Danaus plexippus, 682
endangered biological phenomenon, 682
North America, 682
seasonal migration, 682
population distribution pattern,
Ceratitis capitata, 77
invasion, 77
overwintering, 77
spatial patterns, 77
precocene,
Drosophila melanogaster, 601
female receptivity, 601
ovaries, 601
oviposition, 601
predation,
Aphis gossypii, 475
Lemnia biplagiata, 475
Life table, 475
prediapause,
diapause, 575
Dolycoris baccarum, 575
fatty acid composition, 575
Piezodorus lituratus, 575
prevalence,
Harmonia axyridis, 96
ladybird beetle, 96
male-killing, 96
Spiroplasma, 96
primary host plant,
Aphis glycines, 926
buckthorn, 926
Rhamnus alnifolia, 926

Rhamnus cathartica, 926
probing,
electrical penetration graph, 787
electronic monitoring of insect feeding, 787
Homalodisca coagulata, 787
Pierce's disease, 787
progressive provisioning,
nymphal independence, 143
parental care, 143
shield bug, 143
propagative bacteria,
Heteroptera, 770
squash bug, 770
transmission, 770
protein baits,
Bactrocera cucurbitae, 402
field cages, 402
methodology, 402
Tephritidae, 402
provisioning,
competition, 134
kleptoparasitism, 134
parental care, 134
risk-sensitive behavior, 134
proximate mechanism,
caste differentiation, 732
Formosan subterranean termite, 732
JH titer, 732
soldier proportion, 732
Pseudacteon,
Argentina, 85
biological control, 85
parasitoid, 85
Solenopsis, 85
Pseudasphondylia elaeocarpi,
Cornus controversa, 259
Elaeocarpus sylvestris, 259
gall midge, 259
Pseudasphondylia kiritanii, 259
Pseudasphondylia kiritanii,
Cornus controversa, 259
Elaeocarpus sylvestris, 259
gall midge, 259
Pseudasphondylia elaeocarpi, 259
pseudo-arrhenotoky,
amplified fragment length polymorphism, 359
Broca, 359
DNA fingerprinting, 359
invasive species, 359
pteridophagy,
Old World climbing fern, 887
Pyraloidea, 887
spiders, 887
tenebrionid beetles, 887
Pterotrogus,
Campephilus principalis, 13
feather mites, 13
new species, 13
pupal color,
conditional strategy, 996
polyphenism, 996
pupation sites, 996
pupal dorsal spines,
dorsal setae on A9, 37
La Selva Biological Station, 37
Moraceae, 37
Tortricidae, 37
pupal weight,
body size, 191
fecundity, 191
oviposition, 191
reproduction, 191
pupation sites,
conditional strategy, 996
polyphenism, 996
pupal color, 996
Pyraloidea,
Old World climbing fern, 887
pteridophagy, 887
spiders, 887
tenebrionid beetles, 887

R

RAPD-PCR,
Anopheles albitarsis s.l., 908
cryptic species, 908
mtDNA *COI* gene, 908
rearing,
Anoplophora glabripennis, 536
artificial diets, 536
iron, 536
recolonization,
Ceratitis capitata, 302
large-scale population dynamics, 302
local extinction, 302
long-distance dispersal, 302
recovery,
Cicindelidae, 552
tiger beetle, 552
translocation, 552
reproduction,
body size, 191
fecundity, 191
oviposition, 191
pupal weight, 191
Reticulitermes, 395, 980
16S, 980
DNA sequence, 395, 980
genetic variation, 395, 980
termite, 395, 980
Rhagoletis indifferens,
bitter cherry, 703
black hawthorn, 703
cascara, 703
cherry laurel, 703
Rhagoletis mendax,
blueberry maggot, 966
diapause, 966
heat stress, 966
Hsp70, 966
Rhagoletotrypeta cubensis n. sp.,
Celtis, 252
host plants, 252
taxonomy, 252
Tephritidae, 252
Rhamnus,
Aphis glycines, 690
buckthorn, 690
soybean aphid winter host, 690
Rhamnus alnifolia,
Aphis glycines, 926
buckthorn, 926
primary host plant, 926
Rhamnus cathartica, 926
Rhamnus cathartica,
Aphis glycines, 926
buckthorn, 926
primary host plant, 926
Rhamnus alnifolia, 926
Rhopalidae,
biology, 515
Cardiospermum halicacabum, 515
Sapindaceae, 515
setal patterns, 515
Rhopalomyia californica,
^{13}C, 205
^{15}N, 205
parasitoids, 205
stable isotopes, 205
Rhyparochromidae,
Heteroptera, 458
Mexico, 458
Myodocha, 458
risk-sensitive behavior,
competition, 134
kleptoparasitism, 134
parental care, 134
provisioning, 134

S

16S,
DNA sequence, 980
genetic variation, 980
Reticulitermes, 980
termite, 980
Sabethini,
Culicidae, 838
new species, 838
systematics, 838
Wyeomyia, 838
Sapindaceae,
biology, 515
Cardiospermum halicacabum, 515
Rhopalidae, 515
setal patterns, 515
Scarabaeidae,
Cetoniinae, 658
Dasyvalgus, 658
Valgini, 658
Valgus, 658
seasonal flight period,
apple, 214
distribution, 214
emergence, 214
hosts, 214
seasonal migration,
Danaus plexippus, 682
endangered biological phenomenon, 682
North America, 682
population census, 682
SEM,
chorion, 346
morphology, 346
terminal filament, 346
Tipula, 346
sensor structure,
infrared radiation receptor, 738
Melanophila, 738
phasic receptor, 738
transduction, 738
setal patterns,
biology, 515
Cardiospermum halicacabum, 515
Rhopalidae, 515
Sapindaceae, 515
sex ratio,
adult emergence, 336
Pimpla turionellae, 336
vitamin E, 336

sexual selection,
genitalia, 527
Insecta, 527
mating isolation, 527
morphometrics, 527
sharpshooter,
biometric data, 322
developmental biology, 322
Homalodisca coagulata, 322
life table parameters, 322
shield bug,
nymphal independence, 143
parental care, 143
progressive provisioning, 143
shore flies,
Diptera, 431
Discomyza, 431
Ephydridae, 431
malacophagy, 431
Sinea spinipes,
descriptions, 18
laboratory rearing, 18
life history, 18
nymphs, 18
social wasps,
attractiveness, 236
pheromones, 236
yellowjackets, 236
sociality,
eusociality, 126
Madagascar, 126
mass provisioning, 126
phylogenetics, 126
soil moisture,
anoxia, 973
Cicindelidae, 973
flooding., 973
tiger beetle, 973
soldier formation,
caste regulation, 340
Coptotermes formosanus, 340
juvenile hormone, 340
soldier proportion,
caste differentiation, 732
Formosan subterranean termite, 732
JH titer, 732
proximate mechanism, 732
Solenopsis,
Argentina, 85
biological control, 85
parasitoid, 85
Pseudacteon, 85
soybean aphid winter host,
Aphis glycines, 690
buckthorn, 690
Rhamnus, 690
Spain,
Cynipidae, 853
Iraella, 853
new species, 853
Papaver, 853
Spathius agrili,
Agrilus planipennis, 636
biocontrol, 636
new species, 636
parasitoid, 636
spatial patterns,
Ceratitis capitata, 77
invasion, 77
overwintering, 77
population distribution pattern, 77
species complex,
Aleyrodidae, 827
cryptic species, 827
genetic variation, 827
virus vector, 827
species richness,
geographic distribution, 694
Miridae, 694
plant bugs host plants, 694
sperm precedence,
insecticide resistance management, 989
Leptinotarsa decemlineata, 989
mating behavior, 989
spermatogenesis,
B chromosomes, 227
biased sex ratio, 227
spiders,
Old World climbing fern, 887
pteridophagy, 887
Pyraloidea, 887
tenebrionid beetles, 887
Spiroplasma,
Harmonia axyridis, 96
ladybird beetle, 96
male-killing, 96
prevalence, 96
Spiroplasma kunkelii,
corn stunt spiroplasma, 820
Dalbulus maidis, 820
immunofluorescence confocal laser scanning microscopy, 820
mollicutes, 820
squash bug,
Heteroptera, 770
propagative bacteria, 770
transmission, 770
stable isotopes,
^{13}C, 205
^{15}N, 205
parasitoids, 205
Rhopalomyia californica, 205
sterile insect technique,
arthropods, 1
disinfestation, 1
irradiation, 1
sterilization, 1
sterilization,
arthropods, 1
disinfestation, 1
irradiation, 1
sterile insect technique, 1
stridulation,
development, 219
ecdysis, 219
Laupala, 219
sugar-fed flies,
lateral neurosecretory cells, 329
liver-fed flies, 329
median neurosecretory cells, 329
midgut hormone, 329
symbiosis,
Fungus-growing ants, 151
mycophagy, 151
origin, 151
systematics, 838, 861
Chagas disease, 861
Culicidae, 838
Linshcosteus karupus, 861
morphology, 861
new species, 838
Sabethini, 838
Triatominae, 861
Wyeomyia, 838
Systena nitentula,
alligator weed, 643
Alternanthera philoxeroides, 643
Alticinae, 643
biological control, 643

T

Tachinidae,
citrus root weevil, 245
Oestrophasia (*Cenosoma*) *sabroskyi*, 245
parasitoid, 245
taxonomy, 55, 187, 252, 273, 291, 587
arboreal termite, 273
Celtis, 252
Chironomidae, 587
cicada, 187
Cladopelma, 587
Costa Rica, 187
distribution, 273
DNA markers, 587
Dohrniphora, 55
egg parasitoid, 291
Fidicina, 187
host plants, 252
Hymenoptera, 291
Insecta, 291
mitochondrial DNA, 587
Neotropical, 55
new species, 187
Phoridae, 55
Rhagoletotrypeta cubensis n. sp., 252
Tephritidae, 252
Trichogrammatidae, 291
Tektin,
Ithomiinae, 873
Lepidoptera, 873
molecular phylogenetics, 873
tenebrionid beetles,
Old World climbing fern, 887
pteridophagy, 887
Pyraloidea, 887
spiders, 887
Tephritidae, 173, 252, 402
Anastrepha, 173
Bactrocera cucurbitae, 402
C3C5 region, 173
CCID, 173
Celtis, 252
field cages, 402
host plants, 252
methodology, 402
period, 173
protein baits, 402
Rhagoletotrypeta cubensis n. sp., 252
taxonomy, 252
terminal filament,
chorion, 346
morphology, 346
SEM, 346
Tipula, 346
termite, 395, 980
16S, 980
DNA sequence, 395, 980
genetic variation, 395, 980
Reticulitermes, 395, 980

Tetranychus kanzawai,
dispersal capacity, 625
mating status, 625
mean crowding, 625
trail, 625
thelytoky,
armored scale insects, 629
Aspidiotini, 629
endosymbiosis, 629
parthenogenesis, 629
thermotolerance,
heat shock proteins, 100
heat treatment, 100
Tribolium castaneum, 100
tiger beetle, 552, 973
anoxia, 973
Cicindelidae, 552, 973
flooding., 973
recovery, 552
soil moisture, 973
translocation, 552
Tipula,
chorion, 346
morphology, 346
SEM, 346
terminal filament, 346
top-down,
bottom-up, 673
coexistence, 673
mosquitoes, 673
treehole, 673
Tortricidae,
dorsal setae on A9, 37
La Selva Biological Station, 37
Moraceae, 37
pupal dorsal spines, 37
trail,
dispersal capacity, 625
mating status, 625
mean crowding, 625
Tetranychus kanzawai, 625
transduction,
infrared radiation receptor, 738
Melanophila, 738
phasic receptor, 738
sensor structure, 738
translocation,
Cicindelidae, 552
recovery, 552
tiger beetle, 552
transmission,
Heteroptera, 770
propagative bacteria, 770
squash bug, 770
treehole,
bottom-up, 673
coexistence, 673
mosquitoes, 673
top-down, 673
Triatominae,
Chagas disease, 861
Linshcosteus karupus, 861
morphology, 861
systematics, 861
Tribolium castaneum,
heat shock proteins, 100
heat treatment, 100
thermotolerance, 100
Trichogrammatidae,
egg parasitoid, 291
Hymenoptera, 291
Insecta, 291
taxonomy, 291

U

ultrastructure,
antenna, 113
coeloconic sensilla, 113
maxillary palp, 113
Megaselia scalaris, 113

V

Valgini,
Cetoniinae, 658
Dasyvalgus, 658
Scarabaeidae, 658
Valgus, 658
Valgus,
Cetoniinae, 658
Dasyvalgus, 658
Scarabaeidae, 658
Valgini, 658
vector,
feeding behavior, 814
plant disease, 814
vector transmission,
Brevicoryne brassicae, 763
CaMV, 763
inoculation, 763
Myzus persicae, 763
vineyard,
biodiversity, 565
Curculionoidea, 565
weevil, 565
virus vector,
Aleyrodidae, 827
cryptic species, 827
genetic variation, 827
species complex, 827
vitamin E,
adult emergence, 336
Pimpla turionellae, 336
sex ratio, 336
volume,
biomass, 548
length–mass relationship, 548
Neotropical Scarabaeidae, 548
volume–mass relationship, 548
volume–mass relationship,
biomass, 548
length–mass relationship, 548
Neotropical Scarabaeidae, 548
volume, 548

W

water hyacinth,
biological control, 66
Eichhornia crassipes, 66
immature stages, 66
Megamelus scutellaris, 66
weevil,
biodiversity, 565
Curculionoidea, 565
vineyard, 565
Wellington Formation,
Coleoptera, 73
fossil insects, 73
Permian, 73
West Indies,
biogeography, 951
Florida, 951
over-water dispersal, 951
peninsula effect, 951
wings,
aculeus, 367
chromosomes, 367
Haldane's rule, 367
isozymes, 367
Wyeomyia,
Culicidae, 838
new species, 838
Sabethini, 838
systematics, 838

Y

yellowjackets,
attractiveness, 236
pheromones, 236
social wasps, 236